



Examination of the Physical Aspects of Hydrogen Storage in MOFs

Omar M. Yaghi and Adam J. Matzger
Department of Chemistry
University of Michigan
Ann Arbor, MI 48109

This presentation does not contain any proprietary or confidential information

Project ID #
STP39 Yaghi



Overview

Timeline

- Project start date
1/1/2005
- Project end date
12/31/2009
- Percent complete
2.5%

Budget

- Total project funding
 - DOE share: \$ 1.75M
 - Contractor share: \$ 0.437M
- Funding received in FY04
 - \$0.00
- Funding for FY05
 - \$62,500

Barriers

- Technical barriers addressed
 - B) Weight and Volume
 - C) Efficiency
 - M) Hydrogen Capacity and Reversibility
 - N) Lack of Understanding of Hydrogen Physisorption and Chemisorption
- Technical targets by YR 2010
 - Gravimetric capacity: 6.0%
 - Volumetric capacity: 4.5%
 - Operating ambient temp.: -30/50 °C
 - Cycle life: 1000


Partners

- NREL Team
- Yue Wu (University of North Carolina)
- Hansong Chen (Air Products)



CbHS Center of Excellence Partners

9 university projects (at 7 universities), 4 government labs, 1 industrial partner





Objectives

To develop novel, highly porous metal-organic framework materials (MOFs) as high capacity sorbents for H₂ storage applications.

- Assess gravimetric and volumetric H₂ storage capacities of MOFs.
 - Explore pressure and temperature dependence of H₂ uptake in existing MOF materials over the parameter range specified in DOE YR2010 guidelines.
- Determine the optimal pore size and functionality for H₂ sorption and release in MOFs.
 - Characterize H₂ adsorption sites in existing MOF materials.
 - Study the relationship between pore size and level of H₂ uptake.



Approach

- Equilibrium H₂ uptake as a function of structure.
 - Survey hydrogen uptake levels in existing MOFs, under a variety conditions, to determine promising materials.
 - Use Raman Spectroscopy to probe in H₂ sorbed in pores.
- Correlate systematic changes in organic links with uptake.
 - Use above results to aid design of new structures with greater hydrogen storage capacities.
 - Synthesize new organic linkers and the corresponding MOFs.
 - Measure equilibrium H₂ uptake.



General Synthetic Strategy of MOFs















Inorganic Units

SBUs

Organic Units


SBUs






Control of MOF Dimensionality


1D



2D




3D








Synthesis of an Isoreticular Series of MOFs





Variation in MOF Metrical Parameters






Mechanism of H₂ Uptake in MOFs

Chemisorption

Mg-Hydride, 300 °C


Weight % H₂



Physisorption


MOF-5, 24 °C

Weight % H₂







Inelastic Neutron Scattering of H₂ in MOF-5



Smallest repeat unit of MOF-5





Extended structure of MOF-5







H₂ Sorption at 1 atm & 77 K



IRMOF-1






IRMOF-18



IRMOF-8





Synthesis of a Cu-based MOF

















MOF-505 H₂ (77 K) Sorption Isotherms– Activation Study



MOF-505



Future Work

TASK	2005	2006	2007	2008	2009	
Task 1: Equilibrium H₂ Uptake vs Structure Measure temperature and pressure dependence of H ₂ uptake in 4 existing MOFs Computational design of new linkers Explore new linkers		  	 Go/ No-go Option	 Go/ No-go Option		
Task 2: Thermodynamics & Kinetics H₂ Uptake Determine H ₂ binding energy to framework by sorption Analysis H ₂ uptake rates in single crystals by Raman spectroscopy Measure H ₂ flux in single x-tals	  					
Task 3: Mechanism of H₂ Uptake Analysis H ₂ binding sites in MOFs by Raman spectroscopy NMR measurement of H ₂ uptake		 				
Task 4: Down-select Optimum Materials Optimize scale-up synthesis and activation Cost analysis of material			 			



Publications and Presentations

Please list any publications and presentations that have resulted from work on this project.

- *No publications resulting from current funding at this stage of the project.*



Hydrogen Safety

The most significant hydrogen hazard associated with this project is:

- High exposure to H₂ gas with possibility of personal injury due to decreased oxygen content in the atmosphere.
- High concentrations of H₂ may pose a fire or explosion in and around instrumentation.



Hydrogen Safety

Our approach to deal with this hazard:

- Dedicated a single laboratory for all H₂ experiments.
- Installed active ventilation snorkles from laboratory hoods to all instrumentation consuming/ releasing H₂.
- Installed atmospheric H₂ detector (ppm level detection) outfitted with an alarm in the dedicated laboratory.



Acknowledgements

